

Fermions in a Multi-Soliton Background in 1+1 Dimensions

B. Altschul¹

*Department of Mathematics
Massachusetts Institute of Technology
Cambridge, MA, 02139-4307 USA*

Abstract

We consider the fermionic bound states associated with a soliton-antisoliton pair in 1+1 dimensions which have zero energy when the solitons are infinitely far apart. We calculate the energies of these states when the solitons are separated by a finite distance. The energies decay exponentially with the distance between the soliton and antisoliton. When the fermion mass is much larger than the boson mass, the energy simplifies substantially. These energies may be interpreted as a contribution to the effective potential between the soliton and antisoliton. The character of this contribution depends upon which fermionic states are occupied. Performing the analogous calculation for the simplest (3+1)-dimensional soliton system, we find no fermionic energy shift.

¹baltschu@mit.edu

The (1+1)-dimensional Dirac equation, when coupled to a scalar field, has the interesting property that with the scalar field in a single-soliton state, the fermionic spectrum can include a single zero-energy mode, so that the electric charge of the system is half-integral [1]. However, while the one-soliton case is exactly solvable, the problem of fermions in the presence of multiple interacting solitons is difficult to analyze exactly. Graham and Jaffe [2] have developed a numerical method for calculating the fermionic energies in a multi-soliton background. We shall examine the two-soliton case using an alternative method, deriving an analytic approximation for the fermionic energies. Our approximation method can also be applied to problems in 3+1 dimensions, where numerical calculations are difficult.

The free Dirac equation in 1+1 dimensions is $i\gamma^\mu\partial_\mu\psi + m\psi = 0$, where a suitable choice for the Dirac matrices is $\alpha = \sigma_2$ and $\beta = \sigma_1$. If we use this representation, then the Hamiltonian $H = \alpha p + \beta m$ becomes

$$H = -i\sigma_2\partial_x + \sigma_1m = \begin{bmatrix} 0 & -\partial_x + m \\ \partial_x + m & 0 \end{bmatrix}. \quad (1)$$

We shall replace the fermion mass m with a scalar potential $U(x)$, which is time-independent (or varies very slowly in time), so that energy is conserved.

We are interested in the case where the potential $U(x)$ that appears in the Dirac equation is derived from a scalar field ϕ . We shall take the Lagrange density for this field to be

$$\mathcal{L} = \frac{1}{2}\dot{\phi}^2 - \frac{1}{2}(\partial_x\phi)^2 - V(\phi). \quad (2)$$

We shall concentrate on the “Mexican hat” ϕ^4 potential,

$$V(\phi) = \frac{\lambda^2}{4} \left(\phi^2 - \frac{a^2}{\lambda^2} \right)^2, \quad (3)$$

although similar results hold for the sine-Gordon potential [3, 4].

The ϕ^4 potential has two degenerate vacua, at $\phi = \pm \frac{a}{\lambda}$. There are also solutions that interpolate between the two vacua (i.e. solitons). For a stationary soliton centered at $x = x_0$, the solution is

$$\phi_{S,x_0}(x) = \frac{a}{\lambda} \tanh \left[\frac{\mu}{2}(x - x_0) \right], \quad (4)$$

where $\mu^2 \equiv 2a^2$ is the mass squared of the light quanta of the scalar field. As $x \rightarrow \pm\infty$, $\phi_{S,x_0}(x) \rightarrow \pm \frac{a}{\lambda}$. There is also an antisoliton solution $\phi_{A,x_0}(x) = -\phi_{S,x_0}(x)$, which goes the other direction, from $+\frac{a}{\lambda}$ at $x = -\infty$ to $-\frac{a}{\lambda}$ at $+\infty$. Finally, there are solutions that resemble spatially alternating solitons and antisolitons. However, treating these solutions as collections of individual solitons and antisolitons is a good approximation only when the solitons and antisolitons are widely separated, so that they disturb one-another only minimally. In general, these solitons must be moving relative to one-another, but we shall

assume that this motion is very slow, so that $\phi(x)$ depends very weakly on time. In three spatial dimensions, there do exist exact multi-soliton (multi-monopole) static solutions to bosonic field equations.

We may introduce massless fermions coupled to this scalar field by a Yukawa interaction, $U(x) = g\phi(x)$. When ϕ is in a vacuum state, this generates a fermion mass $m = g\frac{a}{\lambda}$. When ϕ has the soliton profile ϕ_{S,x_0} , there is known to be a zero-energy bound state localized around the zero of the soliton [1, 4, 5], with wavefunction

$$\psi_+ = C \left\{ \cosh \left[\frac{\mu}{2}(x - x_0) \right] \right\}^{-2m/\mu} \begin{bmatrix} 1 \\ 0 \end{bmatrix}. \quad (5)$$

C is a real normalization constant. For the antisoliton ϕ_{A,x_0} the corresponding zero mode wavefunction is

$$\psi_- = C \left\{ \cosh \left[\frac{\mu}{2}(x - x_0) \right] \right\}^{-2m/\mu} \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \quad (6)$$

We shall investigate how the energies of the fermion bound states are affected in the two-soliton case. We shall consider an antisoliton centered at x_1 and a soliton centered at x_2 , where $\mu(x_2 - x_1) \gg 1$. We shall approximate ϕ as a sum $\phi = \phi_{A,x_1} + \phi_{S,x_2} + \frac{a}{\lambda}$ and use ψ_+ and ψ_- to represent the bound state wavefunctions (5) and (6) centered at x_2 and x_1 , respectively.

When the solitons are far apart, we may expect the potential due to one soliton to be a small perturbation of the Dirac equation for the bound state localized around the other soliton. In this spirit, we shall calculate the energy expectation values for the symmetrized states $\Psi_{\pm} \equiv \frac{1}{\sqrt{2}}(\psi_+ \pm \psi_-)$. For $H\psi_+$, we have

$$\begin{aligned} H\psi_+ &= [-i\sigma_2\partial_x + \sigma_1 g\phi_{S,x_2}]\psi_+ + \sigma_1 g \left[\phi_{A,x_1} + \frac{a}{\lambda} \right] \psi_+ \\ &= 0 + \sigma_1 [g\phi_{A,x_1} + m]\psi_+. \end{aligned} \quad (7)$$

We shall first evaluate the energy expectation values $E_{\pm} \equiv \int_{-\infty}^{\infty} dx \Psi_{\pm}^{\dagger} H \Psi_{\pm}$ under the approximations $\frac{\mu}{m} \gg 1$ and $\mu(x_2 - x_1) \gg 1$, where elementary expressions can be obtained explicitly. Then we shall present an exact analysis for general $\frac{\mu}{m}$, which reduces to the previous when $\frac{\mu}{m} \gg 1$.

To find $E_{\pm}$, we need to integrate quantities such as $(\psi_+^{\dagger} \sigma_1 \psi_-) \phi_{A,x_1}$. The characteristic size of ϕ_{A,x_1} is $\frac{1}{\mu}$, and the characteristic decay length of $\psi_{\pm}$ is $\frac{1}{m}$. If $\frac{\mu}{m} \gg 1$, then the fermion wavefunctions decay very little over the width of the antisoliton, and it is a good approximation to replace ϕ_{A,x_1} by the step function $-\frac{a}{\lambda} \text{sgn}(x - x_1)$ in the integral. Approximating ϕ_{A,x_1} with this step function gives

$$H\psi_+ = \sigma_1 m [1 - \text{sgn}(x - x_1)] \psi_+. \quad (8)$$

The corresponding expression for $H\psi_-$ is

$$H\psi_- = \sigma_1 m [1 + \text{sgn}(x - x_2)] \psi_-. \quad (9)$$

Combining (8) and (9), we get

$$H\Psi_{\pm} = \frac{1}{\sqrt{2}}\sigma_1 m \{ [1 - \text{sgn}(x - x_1)]\psi_+ \pm [1 + \text{sgn}(x - x_2)]\psi_- \}. \quad (10)$$

Since $\psi_+^\dagger \sigma_1 \psi_+ = \psi_-^\dagger \sigma_1 \psi_- = 0$, only the cross terms in $\Psi_{\pm}^\dagger H\Psi_{\pm}$ are nonzero. Using the relations

$$\psi_+^\dagger \sigma_1 \psi_- = \psi_-^\dagger \sigma_1 \psi_+ = C^2 \left\{ \cosh \left[\frac{\mu}{2}(x - x_1) \right] \cosh \left[\frac{\mu}{2}(x - x_2) \right] \right\}^{-2m/\mu}, \quad (11)$$

the expression for $E_{\pm}$ is

$$\begin{aligned} E_{\pm} = & \pm m \frac{C^2}{2} \int_{-\infty}^{\infty} dx \left\{ \cosh \left[\frac{\mu}{2} \left(x - y + \frac{\Delta x}{2} \right) \right] \cosh \left[\frac{\mu}{2} \left(x - y - \frac{\Delta x}{2} \right) \right] \right\}^{-2m/\mu} \\ & \times \left[2 - \text{sgn} \left(x - y + \frac{\Delta x}{2} \right) + \text{sgn} \left(x - y - \frac{\Delta x}{2} \right) \right], \end{aligned} \quad (12)$$

where $y \equiv \frac{x_2 + x_1}{2}$, and $\Delta x \equiv x_2 - x_1$ is the separation between the solitons. Shifting the integration $x \rightarrow (x - y)$ gives

$$\begin{aligned} E_{\pm} = & \pm m \frac{C^2}{2} \int_{-\infty}^{\infty} dx \left\{ \cosh \left[\frac{\mu}{2} \left(x + \frac{\Delta x}{2} \right) \right] \cosh \left[\frac{\mu}{2} \left(x - \frac{\Delta x}{2} \right) \right] \right\}^{-2m/\mu} \\ & \times \left[2 - \text{sgn} \left(x + \frac{\Delta x}{2} \right) + \text{sgn} \left(x - \frac{\Delta x}{2} \right) \right], \end{aligned} \quad (13)$$

which depends only on Δx . Thus we see that the zero-energy eigenstate of the one-soliton background bifurcates into two states, symmetrically displaced above and below $E = 0$. To proceed further with our estimate of $E_{\pm}$, we must approximate the wavefunctions, using

$$\cosh \left[\frac{\mu}{2}(x - x_0) \right] \approx \frac{1}{2} \left[\theta(x - x_0) e^{\frac{\mu}{2}(x - x_0)} + \theta(x_0 - x) e^{-\frac{\mu}{2}(x - x_0)} \right]. \quad (14)$$

The approximation (14) is natural in this context, since the approximate wavefunctions produced by (14) are the exact wavefunctions corresponding to perfectly rectangular solitons. Using (14), $E_{\pm}$ reduces to an integral over exponentials and step functions, which may be evaluated exactly, yielding,

$$E_{\pm} \approx \pm (16)^{m/\mu} C^2 e^{-m\Delta x}. \quad (15)$$

It only remains to calculate C . Using the same approximation (14) for cosh and demanding that $\int \psi_+^\dagger \psi_+ dx = 1$ gives

$$C^2 \approx (16)^{-m/\mu} m, \quad (16)$$

so that

$$E_{\pm} \approx \pm m e^{-m\Delta x}. \quad (17)$$

An interaction energy of this nature has been previously suggested [1].

We are now in a position to show that the $\Psi_{\pm}$ are the correct approximate states for which to calculate the energy. If we instead used some other linear combination ($b\psi_+ + c\psi_-$), with $|b|^2 + |c|^2 = 1$, the expectation value of the energy would be $E' = 2(\text{Re}\{bc^*\})E_+$. E' takes its minimum value (corresponding to the best approximation to the lowest-energy wavefunction) when $c = -b$. The best approximation to the higher-energy wavefunction must be orthogonal to our expression for the lower-lying state, and so has $c = b$. So the best linear combinations of ψ_+ and ψ_- are $\frac{1}{\sqrt{2}}(\psi_+ \pm \psi_-)$.

To obtain the result (17), we needed to make the approximation $\frac{\mu}{m} \gg 1$. We now present an exact evaluation, which shows that the above is the leading term for large $\frac{\mu}{m}$.

If we insert the exact expressions (4), (5), and (6) into (7) (and the corresponding expression for $H\psi_-$), we get

$$\begin{aligned} H\Psi_{\pm} = & \frac{1}{\sqrt{2}}\sigma_1 m \left\{ 1 - \tanh \left[\frac{\mu}{2} \left(x - y + \frac{\Delta x}{2} \right) \right] \right\} \psi_+ \\ & \pm \frac{1}{\sqrt{2}}\sigma_1 m \left\{ 1 + \tanh \left[\frac{\mu}{2} \left(x - y - \frac{\Delta x}{2} \right) \right] \right\} \psi_-. \end{aligned} \quad (18)$$

Multiplying this by $\Psi_{\pm}^{\dagger}$ from the left and integrating over the shifted variable $(x - y)$ gives

$$\begin{aligned} E_{\pm} = & \pm m \frac{C^2}{2} \int_{-\infty}^{\infty} dx \left\{ \text{sech} \left[\frac{\mu}{2} \left(x + \frac{\Delta x}{2} \right) \right] \text{sech} \left[\frac{\mu}{2} \left(x - \frac{\Delta x}{2} \right) \right] \right\}^{2m/\mu} \\ & \times \left\{ 2 - \tanh \left[\frac{\mu}{2} \left(x + \frac{\Delta x}{2} \right) \right] + \tanh \left[\frac{\mu}{2} \left(x - \frac{\Delta x}{2} \right) \right] \right\}. \end{aligned} \quad (19)$$

When the integral is performed, the two tanh terms contribute equally, giving

$$\begin{aligned} E_{\pm} = & \pm m C^2 \int_{-\infty}^{\infty} dx \left\{ \text{sech} \left[\frac{\mu}{2} \left(x + \frac{\Delta x}{2} \right) \right] \text{sech} \left[\frac{\mu}{2} \left(x - \frac{\Delta x}{2} \right) \right] \right\}^{2m/\mu} \\ & \times \left\{ 1 + \tanh \left[\frac{\mu}{2} \left(x - \frac{\Delta x}{2} \right) \right] \right\}. \end{aligned} \quad (20)$$

We may simplify (20) using hyperbolic identities, in particular

$$\text{sech}(a + b) = \frac{\text{sech } a \text{ sech } b}{1 + \tanh a \tanh b}. \quad (21)$$

Letting $a = \frac{\mu}{2}(x - \frac{\Delta x}{2})$ and $b = \frac{\mu}{2}\Delta x$ in (21) gives

$$E_{\pm} = \pm m C^2 \text{sech}^n \left(\frac{m}{n} \Delta x \right) \int_{-\infty}^{\infty} dx \frac{\text{sech}^{2n} \left[\frac{m}{n} \left(x - \frac{\Delta x}{2} \right) \right] \left\{ 1 + \tanh \left[\frac{m}{n} \left(x - \frac{\Delta x}{2} \right) \right] \right\}}{\left\{ 1 + \alpha \tanh \left[\frac{m}{n} \left(x - \frac{\Delta x}{2} \right) \right] \right\}^n}, \quad (22)$$

where we have introduced the dimensionless parameters $n \equiv \frac{2m}{\mu}$ and $\alpha \equiv \tanh(\frac{m}{2}\Delta x)$. Making the substitution $u = \tanh[\frac{m}{n}(x - \frac{\Delta x}{2})]$ in (22) simplifies this integral substantially, giving

$$E_{\pm} = \pm m C^2 \operatorname{sech}^n \left(\frac{m}{n} \Delta x \right) \int_{-1}^1 du \frac{n}{m} \frac{(1+u)^n (1-u)^{n-1}}{(1+\alpha u)^n}. \quad (23)$$

To get a final expression for the energy, we must again calculate C . The normalization gives

$$\begin{aligned} \frac{1}{C^2} &= \int_{-\infty}^{\infty} dx \operatorname{sech}^{2n} \left[\frac{m}{n} (x - x_0) \right] \\ &= \int_{-1}^1 du \frac{n}{m} (1-u^2)^{n-1} \\ &= \frac{n}{m} \sqrt{\pi} \frac{\Gamma(n)}{\Gamma(n + \frac{1}{2})}, \end{aligned} \quad (24)$$

which agrees with (16) as $n \rightarrow 0$.

The remaining integral in (23) may also be found analytically. The result involves the hypergeometric function $F(a_1, a_2; b_1; z)$. In terms of this F , the integral is

$$\int_{-1}^1 du \frac{(1+u)^n (1-u)^{n-1}}{(1+\alpha u)^n} = \sqrt{\pi} \frac{\Gamma(n)}{\Gamma(n + \frac{1}{2})} (1+\alpha)^{-n} F \left(n, n; 1+2n; \frac{2\alpha}{1+\alpha} \right). \quad (25)$$

Fortunately, this simplifies significantly when combined with (24), so that the final formula for the fermion energies is

$$E_{\pm} = \pm m \operatorname{sech}^n \left(\frac{m}{n} \Delta x \right) (1+\alpha)^{-n} F \left(n, n; 1+2n; \frac{2\alpha}{1+\alpha} \right). \quad (26)$$

Since the solitons are far apart, $\alpha \approx 1 - 2e^{-\frac{2m}{n}\Delta x} \equiv 1 - \epsilon$ is close to one, and it is natural to expand (25) around $\alpha = 1$. If we expand the integral in (25) to $\mathcal{O}(\epsilon)$, we find,

$$\int_{-1}^1 du \frac{(1+u)^n (1-u)^{n-1}}{(1+\alpha u)^n} = \int_{-1}^1 du (1-u)^{n-1} + \epsilon \left[n \int_{-1}^1 du \frac{u(1-u)^{n-1}}{(1+u)} \right]. \quad (27)$$

The $\mathcal{O}(\epsilon^0)$ term in (27) has value $\frac{2^n}{n}$. However, the $\mathcal{O}(\epsilon)$ term is divergent, because $F(n, n; 1+2n; z)$ is not analytic at $z = 1$. Instead, we have

$$\int_{-1}^1 du \frac{(1+u)^n (1-u)^{n-1}}{(1+\alpha u)^n} = \frac{2^n}{n} + n 2^{n-1} (\epsilon \ln \epsilon) + \mathcal{O}(\epsilon). \quad (28)$$

Using (28) and $\epsilon = 2e^{-\frac{2m}{n}\Delta x}$, the $\mathcal{O}(\epsilon \ln \epsilon)$ expression for $E_{\pm}$ is

$$E_{\pm} \approx \pm m \left[\frac{2^n}{\sqrt{\pi}} \frac{\Gamma(n + \frac{1}{2})}{\Gamma(n+1)} \right] \operatorname{sech}^n \left(\frac{m}{n} \Delta x \right) \left[1 - n (2m \Delta x) e^{-\frac{2m}{n}\Delta x} \right]. \quad (29)$$

As $n \rightarrow 0$, this agrees with the earlier result (17), through terms of $\mathcal{O}(n)$.

The integral (25) may be evaluated in terms of elementary functions whenever n is an integer. Of particular interest is the case $n = 2$, which corresponds to a supersymmetric Lagrangian. For that case, the integral is

$$\int_{-1}^1 du \frac{(1+u)^2(1-u)}{(1+\alpha u)^2} = \frac{1}{\alpha^4} \left[(-4\alpha^2 + 6\alpha) + (\alpha^2 + 2\alpha - 3) \ln \left(\frac{1+\alpha}{1-\alpha} \right) \right], \quad (30)$$

and the energy is

$$E_{\pm} = \pm \frac{3}{2} m \operatorname{sech}^2 \left(\frac{m}{2} \Delta x \right) \left\{ \frac{1}{\alpha^4} \left[(-2\alpha^2 + 3\alpha) + \left(\frac{m}{2} \Delta x \right) (\alpha^2 + 2\alpha - 3) \right] \right\}. \quad (31)$$

The first term on the right-hand-side of (30) gives the $\mathcal{O}(\epsilon^0)$ term, while the second term is $\mathcal{O}(\epsilon \ln \epsilon)$.

The expression (26) gives the energies for the fermion bound states that lie closest to zero energy. When $n < 1$, the one-soliton system has only the single bound state (5) [5], so the soliton-antisoliton system we are considering has two bound states, with energies given by (26). The fermion energies depend on Δx , so (26) generates a contribution to the effective potential between the soliton and the antisoliton. The character of this potential term depends on which fermionic states are occupied. In the vacuum, the state Ψ_- is occupied, while Ψ_+ is empty. E_- is negative and for $\alpha \approx 1$ becomes more negative as Δx decreases, creating an attractive interaction between the soliton-antisoliton pair. Since this interaction exists in the absence of any fermions or antifermions, it must be generated by virtual particle effects. If either a single fermion or a single antifermion is present (i.e. both states are either filled or empty), the total energy vanishes, and these states do not contribute to the effective potential. Evidently, the presence of a single particle causes a cancellation of the virtual particle corrections mentioned above. In the presence of both a fermion and an antifermion (Ψ_+ occupied and Ψ_- empty), the potential term is repulsive. This presents an interesting picture, with the fermions introducing novel interactions between the soliton-antisoliton pair.

If the positions of the soliton and antisoliton are reversed, so that $x_1 > x_2$, a calculation of the energies proceeds along the same lines. We could extend (26) by replacing Δx by $|\Delta x|$ (both in the sech term and in α) and multiplying by $\operatorname{sgn}(x_2 - x_1)$. The energies obtained would be valid whenever $\mu|\Delta x| \gg 1$. However, there are additional subtleties that arise when we allow Δx to change signs. This involves the solitons passing through one-another. The ϕ^4 solitons do not pass through one-another to emerge undistorted [6]; a sine-Gordon soliton and antisoliton will pass through each other without distortion, but when the solitons are in precisely the same place, the field takes a vacuum value everywhere, so there are no bound states at that instant [3]. In either case, it is impossible to identify which bound state for $x_2 > x_1$ corresponds to which bound state for $x_1 > x_2$. However, (26) does have the attractive feature that the energies $E_{\pm}$ approach the energies

of stationary continuum solutions as $\Delta x \rightarrow 0$ (even though (26) is not valid for small values of the separation).

The simplest (3+1)-dimensional soliton with a known zero-energy fermion mode is the 't Hooft-Polyakov monopole [1, 7] coupled to isospinor fermions (in the fundamental representation). The monopole is described by a scalar field $\Phi_M = \Phi_M^a T^a$ and a vector field $A_M^0 = 0$, $A_M^i = A_M^{ia} T^a$, where T^a is the 2×2 isospin generator. Using the representation

$$\boldsymbol{\alpha} = \begin{bmatrix} 0 & \boldsymbol{\sigma} \\ \boldsymbol{\sigma} & 0 \end{bmatrix}, \quad \beta = i \begin{bmatrix} 0 & -I \\ I & 0 \end{bmatrix} \quad (32)$$

for the Dirac matrices, the associated zero mode is

$$\psi_M = \begin{bmatrix} \chi_+ \\ 0 \end{bmatrix}, \quad \chi_+^{\nu n} = f(|\mathbf{r} - \mathbf{r}_M|)(s_+^\nu s_-^n - s_-^\nu s_+^n), \quad (33)$$

where ν is the spinor index, n is the isospin index, $\mathbf{r}_M$ is the position of the monopole, and f is a known function of the boson fields. The $s_\pm^{\nu n}$ are orthonormal spinor, isospinor basis vectors. The corresponding zero mode for an antisoliton with boson fields Φ_A , $A_A^0 = 0$, and A_A^i is

$$\psi_A = \begin{bmatrix} 0 \\ \chi_- \end{bmatrix}, \quad \chi_-^{\nu n} = f(|\mathbf{r} - \mathbf{r}_A|)(s_+^\nu s_-^n - s_-^\nu s_+^n). \quad (34)$$

To apply our method to this system, we consider the approximate monopole-antimonopole configuration $\Phi = \Phi_M + \Phi_A + \Phi_0$ (where Φ_0 is the vacuum value of the field as $r \rightarrow \infty$), $A^\mu = A_M^\mu + A_A^\mu$ and calculate the expectation value of the energy for the state

$$\Psi_\pm \equiv \frac{1}{\sqrt{2}}(\psi_M + \psi_A) = \frac{1}{\sqrt{2}} \begin{bmatrix} \chi_+ \\ \chi_- \end{bmatrix} \quad (35)$$

in this background. This energy is

$$E_\pm = \frac{1}{2} \int d^3r [\chi_+^\dagger \chi_-^\dagger] \begin{bmatrix} 0 & \boldsymbol{\sigma} \cdot \mathbf{p} - \boldsymbol{\sigma} \cdot \mathbf{A}^a T^a - iG\Phi^a T^a \\ \boldsymbol{\sigma} \cdot \mathbf{p} - \boldsymbol{\sigma} \cdot \mathbf{A}^a T^a + iG\Phi^a T^a & 0 \end{bmatrix} \begin{bmatrix} \chi_+ \\ \chi_- \end{bmatrix}, \quad (36)$$

where G is the strength of the Yukawa coupling. Since the states $\chi_\pm$ are singlets of spin plus isospin, the terms with $\mathbf{p}$ and Φ vanish immediately. The vector field term gives a contribution proportional to $\delta_i^a A^{ia}$, which vanishes for the monopole-antimonopole profile, because A^{ia} is traceless in (i, a) , since it involves ϵ^{iaj} . So in this approximation, there are two degenerate zero-energy fermion modes for the monopole-antimonopole system.

The isovector fermion does not obviously produce zero-energy modes. Its properties are under investigation.

Acknowledgments

This problem was mentioned by G. 't Hooft in reference [4]. The author is also grateful to R. Jackiw and R. Jaffe for many helpful discussions. This work is supported in part by funds provided by the U. S. Department of Energy (D.O.E.) under cooperative research agreement DE-FC02-94ER40818.

References

- [1] R. Jackiw, C. Rebbi, Phys. Rev. D **13**, 3398 (1976).
- [2] N. Graham, R.L. Jaffe, Nucl. Phys. B **544**, 432 (1999); N. Graham, R.L. Jaffe, Nucl. Phys. B **549**, 516 (1999).
- [3] R. Rajaraman, *Solitons and Instantons* (Elsevier, Amsterdam, 1982), p. 16–23, 291–298.
- [4] G. 't Hooft, hep-th/0010225.
- [5] R. F. Dashen, B. Hasslacher, A. Neveu, Phys. Rev. D **10**, 4130 (1974).
- [6] D.K. Campbell, J.F. Schonfeld, C.A. Wingate, Physica D **9**, 1 (1983).
- [7] G. 't Hooft, Nucl. Phys. B **79**, 276 (1974); A. M. Polyakov, JETP Letters **20**, 194 (1974).